

Loop Groups with Infinite Dimensional Targets and their Unitary Representations

Karl-Hermann Neeb

Department of Mathematics, FAU Erlangen-Nürnberg

Higher Structures in Topology and Geometry V (Hamburg)
May 27, 2011



Emmy-
Noether-
Zentrum

Outline

- 1 Hilbert Lie groups
- 2 Loop groups
- 3 Bounded and semibounded representations
- 4 Bounded representations of Hilbert–Lie groups
- 5 Semibounded representations of Kac–Moody groups
- 6 Semibounded projective representations of Hilbert–Lie groups
- 7 Semibounded projective representations of Kac–Moody groups
- 8 Conclusion

1.1. From compact groups to Hilbert–Lie groups

Definition (Hilbert–Lie algebra)

A **Hilbert–Lie algebra** is a Lie algebra \mathfrak{k} which is a real Hilbert space whose scalar product is invariant: $([x, y], z) = (x, [y, z])$.

A Lie group K is a **Hilbert–Lie group** if $\mathbf{L}(K) = \mathfrak{k}$ is a Hilbert–Lie algebra.

Finite dimensional Hilbert–Lie algebras are the **compact** Lie algebras.

Theorem (Schue, 1960/61; Structure of Hilbert–Lie algebras)

\mathfrak{k} is an orthogonal direct sum $\mathfrak{k} = \mathfrak{z}(\mathfrak{k}) \oplus \widehat{\bigoplus}_{j \in J} \mathfrak{k}_j$, where \mathfrak{k}_j is simple. If \mathfrak{k} is inf. dim. simple, then $\mathfrak{k} \cong \mathfrak{u}_2(\mathcal{H})$ (*skew-herm. Hilbert–Schmidt ops*) for a Hilbert space \mathcal{H} over \mathbb{R} , \mathbb{C} or \mathbb{H} with $(x, y) = \operatorname{tr}_{\mathbb{R}}(xy^*) = -\operatorname{tr}_{\mathbb{R}}(yx)$.

Example

$\mathbf{U}_2(\mathcal{H}) = \{g \in \mathbf{U}(\mathcal{H}) : \|\mathbf{1} - g\|_2 < \infty\}$ is a Hilbert–Lie group with Lie algebra $\mathbf{L}(\mathbf{U}_2(\mathcal{H})) = \mathfrak{u}_2(\mathcal{H})$. Here $\|X\|_2 = \sqrt{\operatorname{tr}(X^*X)}$.

1.2. Root data of simple Hilbert–Lie algebras

\mathfrak{k} simple Hilbert–Lie algebra

$\mathfrak{t} \subseteq \mathfrak{k}$ maximal abelian (**Cartan subalgebra**), $\mathfrak{t} \cong \ell^2(J, \mathbb{R})$

$\mathfrak{k}_{\mathbb{C}} = \mathfrak{t}_{\mathbb{C}} \oplus \widehat{\bigoplus}_{\alpha \in \Delta} \mathfrak{k}_{\mathbb{C}}^{\alpha}$ (root decomposition), orthogonal direct sum

$\Delta = \Delta(\mathfrak{k}, \mathfrak{t})$ is a **locally finite root system**.

Theorem (Stumme '99, Classif. of infinite locally finite root systems)

$$\begin{aligned} A_J &= \{\varepsilon_i - \varepsilon_j : i \neq j \in J\}, & B_J &= \{\pm\varepsilon_i, \pm\varepsilon_i \pm \varepsilon_j : i \neq j \in J\} \\ C_J &= \{\pm 2\varepsilon_i, \pm\varepsilon_i \pm \varepsilon_j : i \neq j \in J\}, & D_J &= \{\pm\varepsilon_i \pm \varepsilon_j : i \neq j \in J\}. \end{aligned}$$

\Rightarrow **4 iso-classes** of pairs $(\mathfrak{k}, \mathfrak{t})$ (for each cardinality $|J|$):

$$A_J: \quad \mathbb{K} = \mathbb{C}, \quad \mathfrak{k} = \mathfrak{u}_2(\mathcal{H})$$

$$B_J, D_J: \quad \mathbb{K} = \mathbb{R}, \quad \mathfrak{k} = \mathfrak{u}_2(\mathcal{H}) =: \mathfrak{o}_2(\mathcal{H}), \quad \dim(\ker(\mathfrak{t})) \in \{1, 0\}$$

$$C_J: \quad \mathbb{K} = \mathbb{H}, \quad \mathfrak{k} = \mathfrak{u}_2(\mathcal{H}) =: \mathfrak{sp}_2(\mathcal{H}).$$

$\mathfrak{k} = \mathfrak{o}_2(\mathcal{H})$ has **two conjugacy classes** of Cartan subalgebras under $\text{Aut}(\mathfrak{k})$.

2.1. Loop groups and twisted loop groups

Definition (Twisted loop groups)

For a Hilbert–Lie group K ,

$$\mathcal{L}(K) := \{f \in C^\infty(\mathbb{R}, K) : (\forall t \in \mathbb{R}) f(t + 2\pi) = f(t)\}$$

is called the corresponding **loop group**. For an automorphism $\varphi \in \text{Aut}(K)$,

$$\mathcal{L}_\varphi(K) := \{f \in C^\infty(\mathbb{R}, K) : (\forall t \in \mathbb{R}) f(t + 2\pi) = \varphi^{-1}(f(t))\}$$

is called the corresponding **twisted loop group**.

$\mathcal{L}_\varphi(K)$ is a **Fréchet–Lie group** with Lie algebra

$$\mathcal{L}_\varphi(\mathfrak{k}) := \{\xi \in C^\infty(\mathbb{R}, \mathfrak{k}) : (\forall t \in \mathbb{R}) \xi(t + 2\pi) = \mathbf{L}(\varphi)^{-1}\xi(t)\}.$$

Note: $\mathcal{L}_\varphi(K)$ is the group of smooth sections of a K -Lie group bundle $\mathcal{K} = (\mathbb{R} \times K)/\sim$ over $S^1 \cong \mathbb{R}/2\pi\mathbb{Z}$, where $(t + 2\pi, k) \sim (t, \varphi(k))$.

2.2. Loop groups with Hilbert targets

K -Lie group bundles over \mathbb{S}^1 correspond to $\text{Aut}(K)$ principal bundles, hence are classified by

$$\pi_0(\text{Aut}(K))/\text{conj.}$$

Note: $\text{Aut}(K) \cong \text{Aut}(\mathfrak{k})$ if K is 1-connected.

Theorem

The *automorphism groups* of the infinite dimensional simple Hilbert–Lie algebras are given by the *connected* groups

$$\text{Aut}(\mathfrak{o}_2(\mathcal{H})) \cong \text{O}(\mathcal{H})/\{\pm \mathbf{1}\}, \quad \text{Aut}(\mathfrak{sp}_2(\mathcal{H})) \cong \text{Sp}(\mathcal{H})/\{\pm \mathbf{1}\}$$

(*real and quaternionic case*) and the *2-component* group (*complex case*)

$$\text{Aut}(\mathfrak{u}_2(\mathcal{H})) = \text{PU}(\mathcal{H}) \rtimes \{\mathbf{1}, \sigma\}, \quad \sigma: \mathcal{H} \rightarrow \mathcal{H} \text{ antilin. isom. involution.}$$

We thus obtain **4 iso-classes** of twisted loop algebras

$$\mathcal{L}(\mathfrak{o}_2(\mathcal{H})), \quad \mathcal{L}(\mathfrak{u}_2(\mathcal{H})), \quad \mathcal{L}(\mathfrak{sp}_2(\mathcal{H})) \quad \text{and} \quad \mathcal{L}_\sigma(\mathfrak{u}_2(\mathcal{H})).$$

2.3. Double extensions

Definition (Double extensions)

For a Lie algebra \mathfrak{g} with invariant symmetric bilinear form κ (quadratic Lie algebra) and a κ -skew-symmetric derivation D on \mathfrak{g} , the corresponding double extension is the quadratic Lie algebra $(\widehat{\mathfrak{g}}, \widehat{\kappa})$, where

$$\widehat{\mathfrak{g}} := \mathbb{R} \times \mathfrak{g} \times \mathbb{R}$$

$$[(z_1, x_1, t_1), (z_2, x_2, t_2)] := (\kappa(Dx_1, x_2), t_1 Dx_2 - t_2 Dx_1 + [x_1, x_2], 0)$$

$$\widehat{\kappa}((z_1, x_1, t_1), (z_2, x_2, t_2)) := z_1 t_2 + z_2 t_1 + \kappa(x_1, x_2).$$

Note: $\widetilde{\mathfrak{g}} := \mathbb{R} \times \mathfrak{g}$ is a central ext. with cocycle $\omega_D(x_1, x_2) := \kappa(Dx_1, x_2)$.
 $\widehat{\mathfrak{g}} \cong \widetilde{\mathfrak{g}} \rtimes_{\widetilde{D}} \mathbb{R}$ for $\widetilde{D}(z, x) := (0, Dx)$.

Ex: $\mathfrak{g} = \mathfrak{u}_2(\mathcal{H})$, $Dx = [T, x]$, $T \in \mathfrak{u}(\mathcal{H})$, $\kappa(Dx, y) = -\text{tr}(T[x, y])$

Rem: \mathfrak{k} Hilbert-Lie algebra \Rightarrow Any 2-cocycle ω can be written as $\omega(x, y) = (Dx, y)$ with $D \in \text{der}(\mathfrak{k}) \Rightarrow$ double extension $\widehat{\mathfrak{k}}_D$.

2.4. Affine Kac–Moody groups

If \mathfrak{k} is a Hilbert–Lie algebra and $\varphi \in \text{Aut}(\mathfrak{k})$, then the loop algebra $\mathcal{L}_\varphi(\mathfrak{k})$ carries the scalar product $(\xi, \eta) := \int_0^{2\pi} (\xi(t), \eta(t)) dt$ and the derivation $D\xi = \xi'$ is skew-symmetric. This leads to the double extension

$$\widehat{\mathcal{L}}_\varphi(\mathfrak{k}) = \mathbb{R} \oplus \mathcal{L}_\varphi(\mathfrak{k}) \oplus \mathbb{R}$$

$$[(z_1, \xi_1, t_1), (z_2, \xi_2, t_2)] := ((\xi'_1, \xi_2), t_1\xi'_2 - t_2\xi'_1 + [\xi_1, \xi_2], 0)$$

Theorem (N., '02, N./Wockel '09; Integrability Theorem)

If \mathfrak{k} is simple, then there exists a simply connected *Fréchet–Lie group* $\widehat{\mathcal{L}}_\varphi(K)$ with Lie algebra $\widehat{\mathcal{L}}_\varphi(\mathfrak{k})$ and center \mathbb{T} .

Definition

We call $\widehat{\mathcal{L}}_\varphi(K)$ the corresponding *(affine) Kac–Moody group*.

Goal: Understand unitary rep's of $\widehat{\mathcal{L}}_\varphi(K) \Rightarrow$ **We need root data.**

2.5. Root systems for Kac–Moody groups

Here are the candidates for root systems of $\widehat{\mathcal{L}}_\varphi(\mathfrak{k})$:

Theorem (Y. Yoshii, 2006)

The irreducible reduced **locally affine root systems** of **infinite rank** are the following $X_J^{(1)} := X_J \times \mathbb{Z}$ for $X_J \in \{A_J, B_J, C_J, D_J\}$, J an infinite set, and

$B_J^{(2)} := (B_J \times 2\mathbb{Z}) \dot{\cup} ((B_J)_{\text{sh}} \times (2\mathbb{Z} + 1))$, where $(B_J)_{\text{sh}} = \{\pm \varepsilon_j : j \in J\}$.

$C_J^{(2)} := (C_J \times 2\mathbb{Z}) \dot{\cup} (D_J \times (2\mathbb{Z} + 1))$

$BC_J^{(2)} := (B_J \times 2\mathbb{Z}) \dot{\cup} (BC_J \times (2\mathbb{Z} + 1))$, $BC_J := B_J \cup C_J$.

These root systems contain **no root bases** \Rightarrow No Dynkin diagrams.

To obtain root decompositions of $\widehat{\mathcal{L}}_\varphi(\mathfrak{k})$, we **assume**:

\mathfrak{k} is simple, $\mathfrak{t} \subseteq \mathfrak{k}$ maximal abelian,

$\varphi \in \text{Aut}(\mathfrak{k})$ **involution** with $\varphi(\mathfrak{t}) = \mathfrak{t}$ for which

$\mathfrak{t}^\varphi := \{x \in \mathfrak{t} : \varphi(x) = x\} \subseteq \mathfrak{t}^\varphi$ is **maximal abelian**.

Then $\widehat{\mathfrak{t}}^\varphi := \mathbb{R} \times \mathfrak{t}^\varphi \times \mathbb{R} \subseteq \widehat{\mathcal{L}}_\varphi(\mathfrak{k})$ is **maximal abelian**.

For $\alpha: \mathfrak{t} \rightarrow i\mathbb{R}$ and $n \in \mathbb{Z}$ we define $(\alpha, n): \widehat{\mathfrak{t}}^\varphi \rightarrow i\mathbb{R}$ by

$$(\alpha, n)(z, x, t) := \alpha(x) + int.$$

Then the (anisotropic) root system $\widehat{\Delta} := \Delta(\widehat{\mathcal{L}}_\varphi(\mathfrak{k}), \widehat{\mathfrak{t}}^\varphi)$ is

$$\widehat{\Delta} = (\Delta_+ \times 2\mathbb{Z}) \dot{\cup} (\Delta_- \times (2\mathbb{Z} + 1)) \quad \text{with} \quad \Delta_\pm := \Delta(\mathfrak{k}^{\pm\varphi}, \mathfrak{t}^\varphi) \quad \mathfrak{t}^\varphi\text{-weights}.$$

Theorem (Realization of the 7 locally affine root systems)

For $\Delta(\mathfrak{k}, \mathfrak{t}) = X_J$ we obtain $\widehat{\Delta} = \Delta(\widehat{\mathcal{L}}(\mathfrak{k}), \widehat{\mathfrak{t}}) = X_J^{(1)}$,

and $\widehat{\Delta} = \Delta(\widehat{\mathcal{L}}_\varphi(\mathfrak{k}), \widehat{\mathfrak{t}}^\varphi) = X_J^{(2)}$ is obtained for $\varphi(x) = \sigma x \sigma^{-1}$ as follows:

$B_J^{(2)}$ for $\mathfrak{k} = \mathfrak{o}_2(\mathcal{H})$, σ orth. reflection in \mathfrak{t} -inv. hyperplane, \mathfrak{t} type D.

$C_J^{(2)}$ for $\mathfrak{k} = \mathfrak{u}_2(\mathcal{H})$, σ antilinear with $\sigma^2 = -\mathbf{1}$.

$BC_J^{(2)}$ for $\mathfrak{k} = \mathfrak{u}_2(\mathcal{H})$, σ antilinear with $\sigma^2 = \mathbf{1}$ and $\ker(\mathfrak{t}^\varphi) \neq \{0\}$.

Three root systems for $\widehat{\mathcal{L}}(\mathfrak{o}_2(\mathcal{H})) \cong \widehat{\mathcal{L}}_\varphi(\mathfrak{o}_2(\mathcal{H}))$: $B_J^{(1)}$, $D_J^{(1)}$ and $B_J^{(2)}$.

Two root systems for $\widehat{\mathcal{L}}_\sigma(\mathfrak{u}_2(\mathcal{H}))$ (σ antilinear): $C_J^{(2)}$ and $BC_J^{(2)}$.

3.1. Bounded and semibounded representations

Definition

A unitary representation $\pi: G \rightarrow U(\mathcal{H})$ is called **smooth** if the space $\mathcal{H}^\infty := \{v \in \mathcal{H}: G \rightarrow \mathcal{H}, g \mapsto \pi(g)v \text{ smooth}\}$ of smooth vectors is dense.

The derived representation: $d\pi(x)v = \left. \frac{d}{dt} \right|_{t=0} \pi(\exp tx)v, \quad v \in \mathcal{H}^\infty, x \in \mathfrak{g}.$

The support function: $s_\pi: \mathfrak{g} \rightarrow \mathbb{R} \cup \{\infty\}, s_\pi(x) := \sup \text{Spec}(id\pi(x))$

Cone of semiboundedness: $W_\pi := \{x \in \mathfrak{g}: s_\pi \text{ bounded in a nbhd of } x\}.$

Definition

A smooth representation is called **semibounded** if $W_\pi \neq \emptyset$.

It is called **bounded** if $W_\pi = \mathfrak{g}$.

Theorem (N. '08)

π bounded iff $d\pi: \mathfrak{g} \rightarrow \mathfrak{u}(\mathcal{H})$ continuous iff $\pi: G \rightarrow U(\mathcal{H})$ norm-cont.

3.2. Automatic boundedness

Example

If K is **compact**, then every continuous unitary representation is a direct sum of irreducible ones and **irreducible reps are bounded**.

Remark

- (a) If π is semibounded, then W_π is an **open $\text{Ad}(G)$ -invariant convex cone** in \mathfrak{g} .
- (b) If all open invariant cones in $\mathfrak{g}/\mathfrak{z}(\mathfrak{g})$ are trivial, then every **semibounded** irreducible representation of G is **bounded**.
- (c) All open invariant cones in $\mathfrak{g}/\mathfrak{z}(\mathfrak{g})$ are trivial iff all open invariant cones in \mathfrak{g} intersect $\mathfrak{z}(\mathfrak{g})$ (= **fixed points** of $\text{Ad}(G)$).

Examples

Lie algebras \mathfrak{g} for which open invariant cones in $\mathfrak{g}/\mathfrak{z}(\mathfrak{g})$ are trivial:

- (a) **Hilbert–Lie algebras** (**Bruhat–Tits Fixed Point Thm**)
- (b) $\mathfrak{u}(\mathcal{H})$, \mathcal{H} Hilbert space over $\mathbb{R}, \mathbb{C}, \mathbb{H}$.

4. Bounded representations of Hilbert–Lie groups

\mathfrak{k} simple Hilbert–Lie algebra, $\mathfrak{t} \subseteq \mathfrak{k}$ maximal abelian, Δ corresp. roots

Coroots: $\check{\alpha} \in i\mathfrak{t} \cap [\mathfrak{k}_{\mathbb{C}}^{\alpha}, \mathfrak{k}_{\mathbb{C}}^{-\alpha}]$ with $\alpha(\check{\alpha}) = 2$, for $\alpha \in \Delta$

K the 1-connected Lie group with Lie algebra \mathfrak{k} ; $T := \exp(\mathfrak{t})$

$\mathcal{P}_T := \{\lambda \in i\mathfrak{t}' : (\forall \alpha \in \Delta) \lambda(\check{\alpha}) \in \mathbb{Z}\} \cong \text{Hom}(T, \mathbb{T}) \subseteq i\mathfrak{t}'$ (T -weights)

Weyl group: $\mathcal{W} = \langle r_{\alpha} : \alpha \in \Delta \rangle \subseteq \text{GL}(\mathfrak{t}_{\mathbb{C}})$, $r_{\alpha}(x) = x - \alpha(x)\check{\alpha}$.

Theorem (Classification Theorem, N. '98, '11)

*Bounded unitary representations of K are direct sums of irreducible ones.
The irreducible bounded reps π_{λ} are characterized by their T -weight set*

$$\text{conv}(\mathcal{W}\lambda) \cap (\lambda + \mathcal{Q}), \quad \mathcal{Q} = \langle \Delta \rangle_{\text{grp}} \subseteq \mathcal{P}_T \text{ (root group)}.$$

Classification of bounded irreps: $\mathcal{P}_T/\mathcal{W}$ (every $\lambda \in \mathcal{P}_T$ occurs).

Remark

- (a) Bounded reps of K behave like reps of a compact group.
- (b) The continuous representation theory of K is not type I (Boyer '80)

5. Semibounded representations of Kac–Moody groups

$G = \widehat{\mathcal{L}}_\varphi(K)$ (1-connected) as above (7 types), $d := (0, 0, 1) \in \widehat{\mathcal{L}}_\varphi(\mathfrak{k})$

- π irreducible semibounded rep of $G \Rightarrow d \in W_\pi \cup -W_\pi$
(positive/negative energy representations if $\mp id\pi(d)$ bounded below).
- We use that \mathfrak{k}^φ is simple, hence all its open inv. cones are trivial.
- On the minimal/maximal eigenspace of $-id\pi(d)$ we find a bounded irreducible representation ρ_λ of $Z_G(d) \cong \mathbb{T} \times K^\varphi \times \mathbb{R}$

Theorem (Classification Theorem, Part 1)

Irreducible *semibounded* representations π_λ of $\widehat{\mathcal{L}}_\varphi(K)$ are *extremal weight representations* characterized by their $\widehat{\mathfrak{t}}^\varphi$ -weight set

$$\mathcal{P}_\lambda := \text{conv}(\widehat{W}\lambda) \cap (\lambda + \widehat{Q}) \quad \text{with} \quad \text{Ext}(\text{conv}(\mathcal{P}_\lambda)) = \widehat{W}\lambda$$

(\widehat{W} is the Weyl group of $\widehat{\Delta}$). The set of occurring extremal weights λ is

$$\mathcal{P}^\pm := \{\mu \in \mathcal{P}_{\widehat{\mathfrak{T}}^\varphi} : \mp i(\widehat{W}\mu)(d) \text{ bounded from below}\}.$$

Let $\mathcal{P}_d^\pm \subseteq \mathcal{P}^\pm$ denote those elements μ for which $-i\mu(d)$ is minimal/maximal in $\widehat{\mathcal{W}}\mu$. With $c := \mu(-i, 0, 0)$ (**central charge**), the elements $\mu \in \mathcal{P}_d^+$ are characterized by:

$$c \geq 0, \quad |\mu(\check{\alpha})| \leq \frac{2c}{(\alpha, \alpha)}, \quad |\mu(\check{\beta})| \leq \frac{4c}{(\beta, \beta)} \quad \text{for } (\alpha, 1), (\beta, 2) \in \widehat{\Delta}.$$

Theorem (Classification Theorem, Part 2)

Classification of semibounded irreps: $\mathcal{P}^\pm / \widehat{\mathcal{W}} \cong \mathcal{P}_d^\pm / \mathcal{W}$.

Methods:

- **Convex geometry** of $\widehat{\mathcal{W}}$ -orbits (local Coxeter theory).
- **Complex geometry:** Realization of π_λ in holomorphic sections of a complex Hilbert bundle with fiber representation ρ_λ of $Z_G(d)$ over the complex manifold $G/Z_G(d) \cong \mathcal{L}_\varphi(K)/K^\varphi$ (**holomorphic induction**).
- **Harmonic analysis:** Locally defined operator-valued analytic **positive definite functions**; automatic extension.

6. Semibounded projective reps of Hilbert–Lie groups

Problem: Boundedness of representations of a Hilbert–Lie group K is rather restrictive. It excludes important representations like “infinite wedge representations”. These lead to **projective** representations, hence to **central extensions** and further to **double extensions**.

Setup: \mathfrak{k} simple Hilbert–Lie algebra, $\mathfrak{t} \subseteq \mathfrak{k}$ maximal abelian, $\check{\Delta} \subseteq i\mathfrak{t}$ coroots. A **t-invariant continuous cocycle** $\omega(x, y)$ on \mathfrak{k} can be represented by $D \in \text{der}(\mathfrak{k})$ via $\omega(x, y) = (Dx, y)$ and there exists a linear functional $\lambda: \mathfrak{t} \cap [\mathfrak{k}, \mathfrak{k}] \rightarrow i\mathbb{R}$ with

$$\omega(x, y) = i\lambda([x, y]) \quad \text{for } x, y \in \mathfrak{k}.$$

We call λ a **bounded weight** if $\lambda(\check{\alpha}) \in \mathbb{Z}$ for $\alpha \in \Delta$; \mathcal{P}_b set of bd weights.

Definition

For $\lambda \in \mathcal{P}_b$ we write $\widehat{\mathfrak{k}}_\lambda = \mathbb{R} \oplus \mathfrak{k} \oplus \mathbb{R}$ for the corresp. **double extension**.

$\widehat{\mathfrak{t}} := \mathbb{R} \oplus \mathfrak{t} \oplus \mathbb{R} \subseteq \widehat{\mathfrak{k}}_\lambda$ is maximal abelian.

\widehat{K}_λ is the corresponding 1-connected group; $\widehat{T} := \exp \widehat{\mathfrak{t}} \subseteq \widehat{K}_\lambda$.

$\mathcal{P}_{\widehat{T}} \subseteq \text{Hom}(\widehat{\mathfrak{t}}, i\mathbb{R})$ (group of \widehat{T} -weights).

Theorem (Classification Theorem, N. '11)

Irreducible **semibounded** representations π_μ of \widehat{K}_λ are **extremal weight representations** characterized by their $\widehat{\mathfrak{t}}$ -weight set

$$\mathcal{P}_\mu := \text{conv}(\mathcal{W}\mu) \cap (\mu + \mathcal{Q}) \quad \text{with} \quad \text{Ext}(\text{conv}(\mathcal{P}_\mu)) = \mathcal{W}\mu.$$

Put $d := (0, 0, 1) \in \widehat{\mathfrak{t}}$. The **set of occurring extremal weights** is

$$\mathcal{P}^\pm := \{\mu \in \mathcal{P}_{\widehat{\mathfrak{t}}} : \mp i(\mathcal{W}\mu)(d) \text{ bounded from below}\}.$$

By minimizing/maximizing, we get the d -extremal weights

$$\mathcal{P}_d^\pm = \{\mu \in \mathcal{P}_{\widehat{\mathfrak{t}}} : (\forall \alpha \in \Delta) \lambda(\check{\alpha}) > 0 \Rightarrow \pm \mu(\check{\alpha}) \geq 0\}.$$

Classification: $\mathcal{P}^\pm / \mathcal{W} \cong \mathcal{P}_d^\pm / \mathcal{W}_\lambda$, where $\mathcal{W}_\lambda \subseteq \mathcal{W}$ is the stabilizer of λ .

Remark: (a) Representations of \widehat{K}_λ are **projective representations** of K .
(b) For $K = \text{U}_2(\mathcal{H})$ we cover in particular **infinite wedge representations**.

7. Semibounded projective reps of Kac–Moody groups

Again, **projective representations** of $\widehat{\mathcal{L}}_\varphi(K)$ lead to double extensions of $\mathfrak{g} = \widehat{\mathcal{L}}_\varphi(\mathfrak{k})$, hence to **iterated double extensions** $\widehat{\widehat{\mathcal{L}}}_\varphi(\mathfrak{k})$. Here the cocycle is of the form

$$\omega((z_1, \xi_1, t_1), (z_2, \xi_2, t_2)) := \frac{1}{2\pi} \int_0^{2\pi} i\lambda([\xi_1(t), \xi_2(t)]) dt$$

for some bounded weight $\lambda \in \mathcal{P}_b$ for $(\mathfrak{k}, \mathfrak{t})$.

Corresponding Lie groups $\widehat{\widehat{\mathcal{L}}}_\varphi(K)$ exist, and for $d = (0, 0, 1) \in \widehat{\mathcal{L}}_\varphi(\mathfrak{k})$ we have

$$Z_{\mathfrak{g}}(d) = \mathbb{R} \oplus (\mathbb{R} \oplus \mathfrak{k}^\varphi \oplus \mathbb{R}) \oplus \mathbb{R} = \mathbb{R} \oplus \widehat{\mathfrak{k}}_\lambda^\varphi \oplus \mathbb{R}.$$

Semibounded representations of $\widehat{\widehat{\mathcal{L}}}_\varphi(K)$ now lead to **semibounded representations of the double extension** $(\widehat{K}^\varphi)_\lambda$. **These representations are classified!**

Conjecture

Irreducible **semibounded** representations π_μ of $\widehat{\mathcal{L}}_\varphi(K)$ are **extremal weight representations** characterized by their $\widehat{\mathfrak{k}}$ -weight set

$$\mathcal{P}_\mu := \text{conv}(\widehat{\widehat{\mathcal{W}}}\mu) \cap (\mu + \widehat{\widehat{\mathcal{Q}}}) \quad \text{with} \quad \text{Ext}(\text{conv}(\mathcal{P}_\mu)) = \widehat{\widehat{\mathcal{W}}}\mu.$$

The **set of occurring extremal weights** is

$$\mathcal{P}^\pm := \{\mu \in \mathcal{P}_{\widehat{\widehat{\mathcal{T}}}} : \mp i(\widehat{\widehat{\mathcal{W}}}\mu)(d) \text{ bounded from below}\}.$$

By minimizing/maximizing, we get the d -extremal weights \mathcal{P}_d^\pm .

Classification of semibounded irreps: $\mathcal{P}^\pm / \widehat{\widehat{\mathcal{W}}} \cong \mathcal{P}_d^\pm / \widehat{\widehat{\mathcal{W}}}_d$.

Problems: (a) The complex geometric **Banach methods** (holomorphic induction) **break down** because the representations of \widehat{K}_λ are unbounded.

We need a **weaker notion of a complex Hilbert bundle**.

(b) The iterated double extension creates 2 “ d -elements”, but semiboundedness should be controlled by the first one. This requires refined information on **convexity properties of coadjoint orbits**.

Positive energy vs. semiboundedness

- **Semiboundedness is stronger than the positive energy condition** $-id\pi(d)$ **bd. below.** It is crucial that semiboundedness implies boundedness of the K -representation on the minimal energy space. This is automatic if K is compact. In general K has many irreducible unbounded representations which are harder to control, f.i., Boyer's factor representations of $U_2(\mathcal{H})$. We do not expect that the positive energy condition implies semiboundedness in general.
- **Semiboundedness is intrinsic**, it does not refer to the specification of an element $d \in \mathfrak{g}$, such as the positive energy condition. It also does not refer to a specific Cartan subalgebra.
- Our classification results hold for each of the 7 types of root systems of the 4 classes of Lie algebras. For different root systems, resp., conjugacy classes of Cartan subalgebras, we obtain different parameters for the same representations.

Concluding remarks

- **Non-connected loop groups:** $\pi_0(\mathcal{L}(K)) \cong \pi_1(K)$ is non-trivial in general. Which semibounded representations extend to non-connected groups?
- We need a better understanding of the concept of a **Cartan subalgebra** for $\widehat{\mathcal{L}}_\varphi(\mathfrak{k})$. Are there finitely many conjugacy classes?
- Describe the **automorphism group of $\mathcal{L}_\varphi(\mathfrak{k})$** .
- Are there also semibounded representations for double extensions of **mapping groups $C^\infty(M, K)$** , where $\dim M > 1$? The corresponding derivations should correspond to divergence free vector fields on M . Possibly one has to consider **n -fold iterated double extensions**, where $n = \dim M$. Here $M = \mathbb{T}^2$ is the natural testing case.
- For $K = U_2(\mathcal{H})$, \mathcal{H} complex, we have $\text{Aut}(K)_0 \cong \text{PU}(\mathcal{H})$, so that **K -group bundles** with this structure group over X are classified by their **Dixmier–Douady classes** in

$$[X, B\text{PU}(\mathcal{H})] = [X, K(\mathbb{Z}, 3)] \cong H^3(X, \mathbb{Z}).$$